



## Ophthalmic parameters and the presence of clinically significant macular oedema

Knudsen, L.L.; Lervang, H.H.; Lundbye-Christensen, Søren; Gorst-Rasmussen, Anders

*Publication date:*  
2006

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*

Knudsen, L. L., Lervang, H. H., Lundbye-Christensen, S., & Gorst-Rasmussen, A. (2006). *Ophthalmic parameters and the presence of clinically significant macular oedema*. Poster presented at Meeting of the European Association for the Study of Diabetes Eye Complications Study Group (EASDEC), Århus, Denmark.

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -


### Take down policy

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.

# The North Jutland County Diabetic retinopathy Study (NCDRS)

## Retinal lesions and their association to clinically significant macular oedema

LL Knudsen <sup>1A</sup>, HH Lervang <sup>1B</sup>, S Lundbye-Christensen <sup>2</sup> and A Gorst-Rasmussen <sup>2</sup>  
<sup>A</sup>Ophthalmology, <sup>B</sup>Medical endocrinology, <sup>1</sup>Aalborg Sygehus Syd, Århus University Hospital, Aalborg, Denmark; <sup>2</sup>Department of Mathematical Sciences, Aalborg University, Aalborg, Denmark



### Purpose

To explore the influence from retinal lesions on the prevalence of clinically significant macular oedema (CSME) in the present diabetic population.

### Methods


This cross-sectional study comprised 656 type 1 diabetic subjects and 328 type 2 diabetic subjects undergoing retinopathy screening in the County of North Jutland, Denmark during the period 1<sup>st</sup> April 2000 to 30<sup>th</sup> April 2004. The association between CSME and retinal ophthalmic parameters was explored using logistic regression analysis. Type 2 diabetic subjects were recruited from larger Aalborg and comprised more than 75% of known type 1 diabetic subjects in this area. Type 2 diabetic subjects were recruited from the entire county of North Jutland due to malregulation and comprised less than 5% of all known type 2 diabetic subjects.

### Results

The prevalence of proliferative retinopathy was found relatively low in the present diabetic population (Table 1) and lower than previously reported studies (Fig 1). The prevalence of CSME was found relatively high in the present diabetic population (Table 1) and higher than previously reported (Fig 1).


**Table 1**  
Prevalence in NCDRS of proliferative retinopathy and CSME in type 1 and type 2 diabetes.

	Type 1	Type 2
Proliferative retinopathy	0.8 %	0.3 %
CSME	7.9 %	12.8 %




**Fig 1**  
NCDRS and WESDR are compared with respect to HbA1c, blood pressure, prevalence of proliferative retinopathy and CSME. The ratio between medians is depicted, WESCR being reference, i.e. values above one indicates a higher median in the NCDRS.

The prevalence of CSME (Fig 2) and also the number of retinal lesions (Fig 3) revealed a non-linear association to an internationally approved retinopathy scale [1].




**Fig 2**  
The prevalence of CSME at various retinopathy levels.



**Fig 3**  
The number of retinal lesions and their association to an internationally approved retinopathy scale.

The prevalence of CSME increased with the number of retinal lesions (Fig 4) and decreased with improved visual acuity (Fig 4).



**Fig 4**

The association between the various retinal lesions and the presence of CSME are illustrated in Fig 2a (type 1 diabetes) and Fig 2b (type 2 diabetes).

Correlations*						
	Hard exudates	Soft exudates	Intra retinal hemorrhages	Microaneurysms	Visual acuity	CSME
Spearman's rho	1.000	.229**	.350**	.440**	-.104**	.524**
Hard exudates						
Soft exudates	.229**	1.000	.462**	.413**	-.098*	.207**
Intra retinal hemorrhages	.350**	.462**	1.000	.462**	-.176**	.227**
Microaneurysms	.440**	.413**	.462**	1.000	-.110**	.298**
Visual acuity	-.104**	-.098*	-.176**	-.110**	1.000	-.137**
CSME	.524**	.207**	.227**	.298**	-.137**	1.000

\*\* Correlation is significant at the 0.01 level (2-tailed).  
\* Correlation is significant at the 0.05 level (2-tailed).  
a. dmtyp = 1

**Table 2a**  
The association between various retinal lesions and the presence of CSME among type 1 diabetic subjects.

Correlations*						
	Hard exudates	Soft exudates	Intra retinal hemorrhages	Microaneurysms	Visual acuity	CSME
Spearman's rho	1.000	.265**	.589**	.605**	-.166**	.715**
Hard exudates						
Soft exudates	.265**	1.000	.359**	.365**	-.097	.259**
Intra retinal hemorrhages	.589**	.359**	1.000	.707**	-.228**	.585**
Microaneurysms	.605**	.365**	.707**	1.000	-.193**	.508**
Visual acuity	-.166**	-.097	-.228**	-.193**	1.000	-.175**
CSME	.715**	.259**	.585**	.508**	-.175**	1.000

\*\* Correlation is significant at the 0.01 level (2-tailed).  
a. dmtyp = 2

**Table 2b**  
The association between various retinal lesions and the presence of CSME among type 2 diabetic subjects.

## Conclusions

1. The prevalence of CSME seems increased compared to previous studies.
2. The prevalence of CSME is non-linearly associated to an internationally approved retinopathy scale.
3. The prevalence of CSME increased with the number of retinal lesions.